



SOCIETY OF TRAUMA NURSES

THE ELECTRONIC LIBRARY OF

**TRAUMA LECTURES**



THE ELECTRONIC LIBRARY OF

# TRAUMA LECTURES

## Maxillofacial and Ocular Injuries



SOCIETY OF TRAUMA NURSES

# Objectives

**At the conclusion of this presentation  
the participant will be able to:**

- Identify the key anatomical structures of the face and eye and the impact of force on those structures.
- Discuss assessment priorities for a patient with maxillofacial and ocular injuries.
- Prioritize the care of a patient with facial and ocular injuries.
- Discuss psychosocial support for a patient with maxillofacial and ocular injuries.

# Mechanism of Injury

Low  
velocity


High  
velocity





# Pathophysiology


- Bones of face make up the most complex skeletal area of the body.
- Maxillofacial fractures result from either blunt or penetrating trauma.



# Pathophysiology

- 'G' force is a measure of acceleration not produced by gravity
- **High Impact:**
  - Supraorbital rim – 200 G
  - Symphysis Mandible – 100 G
  - Frontal – 100 G
  - Angle mandible – 70 G
- **Low Impact:**
  - Zygoma – 50 G
  - Nasal bone – 30 G







# Etiology

60% of patients with severe facial trauma have multisystem trauma and the potential for airway compromise.








# Etiology

- Approximately one quarter of women with facial trauma are victims of domestic violence.
  - Index of suspicion increases if an orbital wall fx is present.
- Approximately one quarter of patients with severe facial trauma will develop Post Traumatic Stress Disorder.

# Ocular Structures


## Human Eye Anatomy






# Bony Orbit

- Roof
  - Frontal bone
  - Sphenoid
- Medial wall
  - Maxilla
  - Lacrimal, ethmoid
  - Body of sphenoid
- Floor
  - Maxilla
  - Palatine
  - Zygoma
- Lateral
  - Zygoma and greater sphenoid




Je at uwo at English Wikipedia., CC BY 2.5, via Wikimedia Commons

# Cranial Nerves




# Orbital Fractures





# Orbital Fractures


- Usually through floor or medial wall
- Enophthalmos
- Anesthesia
- Diplopia
- Infraorbital stepoff deformity
- Subcutaneous emphysema



Lipa, et al. (2015)

# Orbital Fractures

- Symptoms
  - Periorbital swelling
  - Crepitus
  - Proptosis
  - Ophthalmoplegia
  - Enophthalmos
  - Palpable defects
- Assess for globe injury
- Avoid nose blowing
- Assess for entrapment





Jha, 2018















# Facial Structures

# LeFort I Fracture







# LeFort II Fracture



# LeFort III Fracture




# Le Fort III Fracture


- Periorbital hematoma
- Racoon eyes suggestive of basal skull fracture.
- Inappropriate placement of nasogastric tube.







# Tripod Fracture



# Orbitozygomatic Fractures




- Complex fractures of the zygoma and orbital floor.
- May have double vision, ocular proptosis or enophthalmos.
- Must assess for entrapment of extraocular muscles.
- Surgical management directed at decompression of entrapped muscles and anatomic realignment of zygoma.




# Naso-Ethmoidal-Orbital Fracture


- Fractures that extend into the nose through the ethmoid bones.
- Associated with lacrimal disruption and dural tears.
- Suspect if there is trauma to the nose or medial orbit.
- Patients complain of pain on eye movement.



# Mandibular Fractures







# Mandible Fractures

- Pain
- Malocclusion
- Separation
- Inability to open mouth
- Tongue blade test





# Tongue Blade Test

- Screening test
- Have patient bite tongue blade
- Rotate blade laterally
- 96% sensitive for mandibular fx



Peterson, 2014

# Treatment





# Maxillofacial Injuries General Assessment



- ABC's
- Assess for symmetry of facial structures
  - Assess for paresthesias
  - Assess symmetry of facial movements
- Assess the ears, nose and oral cavity for occult lacerations, hematomas
- Palpate for crepitus, tenderness or deformity
- Assess sense of smell




# Ocular Assessment

- Visual acuity
- Pupil assessment
- Extraocular movements
- Eye position and movement
- Intraocular pressure



# Physical Examination

- Inspect open wounds for foreign bodies.
- Palpate the entire face.
- Inspect the nose.
- Inspect nasal septum for septal hematoma, CSF or blood.
- Palpate nose for crepitus, deformity and subcutaneous air.
- Palpate the zygoma along its arch and its articulations with the maxilla, frontal and temporal bone.







# Physical Examination

- Inspect the teeth
- Intraoral examination:
  - Check for lacerations
  - Stress the mandible
  - Tongue blade test
- Palpate the mandible for tenderness, swelling and step-off





# Physical Examination

- Check visual acuity
- Check pupils for roundness and reactivity
- Examine the eyelids for lacerations
- Test extra ocular muscles
- Palpate around the entire orbits








# Physical Examination

- Examine the cornea for abrasions and lacerations.
- Examine the anterior chamber for blood or hyphema.



# Airway Management

- Protect and maintain airway
  - Pull tongue forward with padded forceps or sutures
  - Endotracheal intubation
  - Anticipate need for cricothyroidotomy
- Prevent aspiration
- Ensure adequate oxygenation and ventilation





# Airway Management

- Protection of airway
- Keep HOB elevated
- Aggressive pulmonary toilet
- Frequent suctioning












# Management

- Control hemorrhage
  - Direct pressure
  - Nasal and oral packing
  - Reduce fractures
- Restore intravascular volume
- Anticipate intracranial injury and need for intervention
  - Serial neurologic exams

# Management

## Nutrition Management

- Early initiation of enteral feeding
- Keep HOB elevated
- Evaluate for swallowing dysfunction prior to oral feeding
- Wire cutters at bedside at all times





# Management


## Prevention of infection

- Perioperative antibiotics
- Frequent oral lavage
- Minimize nasal packing and tubes
- Decongestants
- Avoid blowing nose
- Avoid foreign bodies or instrumentation in nares or ear canal





# Management

- Protect eyes from further injury
- Pain management
- Early Rehab Consult





# Direct Eye Trauma





# Blast Injury: Thermal Injury






# Thermal Injury

- Eye is usually spared
- Corneal exposure may occur as burn heals and skin contracts






# Corneal Abrasion



# Chemical Burns



All Rights Reserved, 2009 Texas Tech University Health Sciences Center EyeAtlas








# Traumatic Hyphema


- Limit activity
- Keep HOB elevated
- Protect the eye
- Monitor intraocular pressure
- Cycloplegic agents
- Monitor for re-bleeding
- Avoid NSAIDs and anticoagulants
- Topical aminocaproic acid

# Lid Laceration

- REFER for
  - Depth
  - Extensive tissue loss
- REFER for location
  - Medial
  - Margin







# Ruptured Globe

- Penetrating
- Blunt
- Urgent ophthalmology consult
- NPO

# Open Globe

- Tetanus
- Antibiotics
- Minimize additional damage
- Avoid increasing intraocular pressure







# Sympathetic Ophthalmia

- Inflammatory condition
- Common after penetrating injury or ruptured globe
- Occurs 5 days to many years after injury
- Results in loss of vision of uninjured eye
- Prevented by early enucleation of injured eye



# Psychosocial Support

- Provide communication aids
- Frequent positive reinforcement
- Early referrals to psychiatric liaisons or counselors
- Early referrals to community agencies for the blind
- Referrals for home safety evaluations
- Referrals to local and state agencies for financial assistance





# Patient and Family Education

- Reinforce surgical plan of care
- Medications
- Nutrition management
- Wound care
- Tracheostomy care
- Avoid direct sunlight for 6-12 months
- Use of cosmetics



# Summary

- Facial and ocular trauma requires a comprehensive multidisciplinary team to maximize outcomes.
- Early incorporation of rehabilitation services is necessary for functional recovery.
- Overall prognosis of reconstruction may take months or years.