



SOCIETY OF TRAUMA NURSES

THE ELECTRONIC LIBRARY OF TRAUMA LECTURES

THE ELECTRONIC LIBRARY OF

TRAUMA LECTURES

Abdominal Trauma





Objectives

**At the conclusion of this presentation
the participant will be able to:**

- Describe common mechanisms of injury seen in abdominal trauma
- Discuss various injuries of the abdomen
- State appropriate assessment and diagnostic studies for the patient with abdominal trauma
- Describe abdominal compartment syndrome and the importance of early recognition



Epidemiology

Incidence

- Abdominal injuries rank among the top seven causes of death in trauma.
- Accounts for more than 10% of trauma deaths
- Seldom a single system injury

Mechanism of Injury

Blunt



Penetrating




By Bobjgalindo - Own work, CC BY-SA 4.0, <https://commons.wikimedia.org/w/index.php?curid=10439138>

Mechanism of Injury

- Heightens suspicion for certain injuries
- Blunt injury and types of forces
- Use of restraint devices
- Penetrating trauma




Anatomy and Physiology



By Tvanbr - Own work, Public Domain,
<https://commons.wikimedia.org/w/index.php?curid=11625998>

Abdominal Sections



Henry Vandyke Carter, Public domain, via Wikimedia Commons




Image courtesy O.Chaigasame (<http://creativecommons.org/licenses/by/4.0/>)

Abdominal Assessment

- Inspection
- Auscultation
- Percussion
- Palpation






Right upper quadrant (RUQ)

Left upper quadrant (LUQ)

Right lower quadrant (RLQ)

Left lower quadrant (LLQ)

Four Quadrants




RUQ

- Liver
- Gallbladder with biliary tree
- Duodenum
- Head of pancreas
- Hepatic flexure of colon

LUQ

- Stomach
- Spleen
- Left lobe liver
- Left kidney
- Left adrenal gland
- Splenic flexure of colon
- Parts of transverse and descending colon





RLQ

- Cecum
- Appendix
- Ascending colon
- Right ovary and fallopian tube
- Right ureter

LLQ

- Descending colon
- Sigmoid colon
- Left ovary and fallopian tube
- Left uterine tube





Ongoing Assessment

- Delayed diagnosis or missed injuries
- Frequent serial and systematic examinations
- Tertiary exam

Diagnostic Labs

Are they necessary?

Reliable?

- Hematocrit
- WBC
- Electrolytes



Diagnostic Labs

- ABGs
- Coagulation studies
- Urinalysis
- Pregnancy
- Pancreatic enzymes
- LFTs



Diagnostic Modalities

- Radiographs
- Diagnostic peritoneal lavage (DPL)
- Ultrasonography (US)
- Computed tomography (CT) scan
- Angiogram
- Diagnostic laparoscopy






Radiographic Films

- CXR
 - Concomitant pulmonary and cardiac injuries
 - Displacement of abdominal organs
- Pelvis
- Plain abdominal films have limited if any role in the acute resuscitation.
 - AP and lateral films may identify fluid or air.
 - Upright film for free air; may disclose ruptured hollow viscus

Diagnostic Peritoneal Lavage

- Used to diagnose intra-abdominal bleeding
- Indications
 - Unexplained hypotension, decreased hematocrit, or shock
 - CT or ultrasound not available
 - Equivocal abdominal examination
 - Altered mental status
 - Spinal cord injury
 - Distracting injuries





DPL

Advantages

- Quick, simple
- Safe
- Low cost
- Relatively accurate
- Grossly positive result

Disadvantages

- Difficult to perform in some patients
- Invasive procedure
- Can miss certain injuries

Complications: Infection, hematoma, false positives, injury, bleeding, unnecessary laparotomy, failure to recover lavage fluid

Note: A urinary catheter and gastric tube should be in place prior to the procedure.

Ultrasound

FAST

- Focused
- Assessment
- Sonography
- Trauma

Ultrasound probe locations and sequence

- Epigastrium
- RUQ
- LUQ
- Pelvis



Ultrasound

- Reliable, fast, safe
- Noninvasive
- Equipment portable
- Performed simultaneously
- Fast exam detects free fluid
- Serial exams
- Leads to fewer DPL's & CT Scans





Disadvantages of Ultrasound

- Clinician expertise variable
- Lacks specificity & sensitivity
- Reliability is questionable
- May not reveal free fluid if performed too early


A Siemens CT scanner is shown in a clinical setting. The large, white, circular gantry is the central feature, with a patient bed positioned inside it. The bed is covered with a white sheet and an orange blanket. The room has a light-colored wall and a wooden door in the background. A trash can is visible on the left side of the frame.

Computed Tomography

- Used for hemodynamically stable patients
- Advantages:
 - Noninvasive procedure
 - Better defines organ injury
 - Estimates amount of blood in spaces
 - Retroperitoneum and vertebrae can be assessed
- Helical scanners

CT Scan in Trauma

- Visualizes abdominal solid organs and vessels well
- Does NOT see mesenteric injuries, hollow viscus, duodenum, diaphragm, or omentum well
- Whole body scans on all trauma
- Radiation long-term effects



CT Scan Disadvantages

- Takes time to perform
- Cost
- Transport of patient
- Requires stable and cooperative patient
- Less reliable in diagnosing some injuries
- IV contrast
- Radiation exposure





Angiography

- Detects active bleeding in patients with vascular trauma
- Embolizes specific structures within bleeding organs or the pelvis
- Detects A-V fistulas and aneurysms in penetrating trauma



Diagnostic Laparoscopy (DL)

- Screening or diagnostic tool
- Invasive procedure with some limitations
- Used to detect or exclude certain findings
- May reduce the rate of negative laparotomies





Other Diagnostic Procedures

ERCP

- May be indicated in the stable trauma patient suspected of having biliary tract or pancreatic duct injury
- Most accurate test in the patient with hyperamylasemia and in those following pancreatic surgery

Other Diagnostic Procedures

Gastrografin or barium studies

- Helpful in diagnosing injuries to the esophagus, stomach, or bowel
- Contrast enemas are used to diagnose rectal or colonic injury secondary to penetrating trauma




Image courtesy S Bhimji MD, <https://www.statpearls.com/ArticleLibrary/viewarticle/21348>

Specific Injuries





Esophageal Injuries



Esophagus

Anatomy

- Carries food from pharynx to the stomach
- Joins the stomach at the level of T-10
- Posterior surface overlies aorta
- Anterior surface covered by peritoneum

Predisposing Injury Facts

- Narrow at
 - Cricoid cartilage
 - Arch of aorta
 - Esophagogastric junction
- Lacks serosal layer
 - Integrity of anastomoses
 - Possible leak after surgical repair



Esophageal Injury

- Incidence
 - Higher in cervical and thoracic areas
 - Majority are due to penetrating trauma
 - Blunt injury is rare
- Early diagnosis essential
- Can result in high morbidity and mortality

Sequelae

- Respiratory compromise
- Mediastinitis
- Paraesophageal abscess
- Empyema
- Esophageal fistula
- Peritonitis



Esophageal Injury

Assessment

- Symptoms of perforation include pain, fever, and dysphagia
- Symptoms of abdominal esophageal tear include signs of peritoneal irritation followed by dyspnea and pleuritic pain

Diagnostic tests

- Endoscopy/Esophagoscopy
- CT

Esophageal Injury




Management

- Initial assessment complex
- Goal is to minimize the bacterial contamination and enzyme erosion
- Gastric decompression
- Antibiotic coverage
- Drainage of wound
- Surgical repair

Esophageal Injury Management

Continuous monitoring for injury

Complications after repair



Samir धर्म, Public domain, via Wikimedia Commons



Diaphragm



Diaphragmatic Injury

Incidence

- Usually occurs with other injuries
- Seen in < 5% of blunt trauma patients
- Left side greater incidence than right side
- Commonly associated with penetrating trauma
- Injuries from blunt trauma caused by sudden rise in intrathoracic pressure



Diaphragmatic Injury


Assessment


- Auscultation of peristaltic sounds in chest
- Delayed rupture - unexplained chest pain and increased respiratory rate
- CXR is most important diagnostic study
 - Elevation of hemidiaphragm
 - Bowel pattern in the chest
 - Gastric tube curls in chest
 - Hemothorax – associated injury
- Masked by positive pressure ventilation



TRAUMA

SUPINE








Diaphragmatic Injury

Management


- Exploratory laparotomy
- Diagnostic laparoscopy in penetrating trauma







Pancreatic Injuries







Pancreatic Injuries

Incidence

- Uncommon (0.2% to 12%)
- Associated with other abdominal injuries
- Majority caused by penetrating trauma
- Blunt trauma is usually a direct blow or compression type force
- Mortality is variable
- Pancreatic Injuries graded by severity I-V
- Most grades of III or higher will require OR



Duodenum



Duodenal Injuries

Incidence


- Majority caused by penetrating trauma
- Blunt trauma is usually compression type
- Mortality is variable
- Multi-organ injuries



Pancreatic and Duodenal Injuries

Assessment

- Peritoneal symptoms not evident but appear later
- CT scan is the exam of choice
- Injury usually found intraoperatively





Pancreatic and Duodenal Injuries

Assessment


- Blunt injury to duodenum can produce intramural hematoma
- Perforation causes contamination



Pancreatic and Duodenal Injuries

Management

- Options depend on site and severity
- Primary closure
- Simple external closed drainage
- Distal pancreatectomy
- Pancreatic duodenectomy





Pancreatic and Duodenal Injuries

Duodenal Management

- Debridement and primary repair
- Surgical procedure depends on hemodynamic stability and duct involvement
- Nonoperative management requires close observation

Pancreatic Management

- Primary cause of death is hemorrhage
- Late deaths are due to sepsis, ARDS, multiple organ failure
- Observe for complications



Injuries to the Stomach and Intestines





Stomach Injury

Incidence

- Rare; more common in children
- Commonly involves adjacent organs
- Protected by location and mobility
- Most common cause is penetrating injury





Stomach Injury

Assessment

- Symptoms variable and nonspecific
- May include severe epigastric or abdominal pain, tenderness, signs of peritonitis
- Clouded by associated injuries
- Bloody output from NG tube
- Free air on radiograph
- Findings on CT or DPL



Stomach Injury

- Indications for surgery
 - Hemodynamic instability
 - Diffuse abdominal pain/peritonitis on clinical exam
 - Studies consistent with GI perforation
- Selective non-operative management
 - Stable
 - No peritoneal signs
 - No clear radiographic evidence




Stomach Injury

Management

- Gastric decompression
- Decision for surgical intervention vs. non-operative management
- If contamination exists, copious peritoneal irrigation and delayed primary closure
- Monitor for postoperative complications

Intestinal Injuries





Small Intestine

Jejunum and Ileum

- Responsible for nutrient absorption and fluid and electrolyte shifts
- Jejunum lies in the umbilical region
- Ileum lies in the hypogastric and pelvic regions
- Vulnerable to seatbelt injury





Small Bowel Injury

Incidence

- Most frequently injured by penetrating trauma.
- Blunt injury is relatively uncommon.
- Presence of pancreatic and solid organ injury are predictive of increased risk for hollow viscus injury.

Assessment

- Clinical signs may not be apparent initially.
- Blunt vs penetrating
- Signs of peritonitis develop.
- Any blow to the abdomen or penetrating injury to the lower chest or abdomen should increase suspicion of injury.




Small Bowel Injury

Management

- Bleeding should be controlled prior to exploration
- Debridement followed by primary closure and ligation of bleeders
- Bowel resection for multiple defects
- Gastric decompression and parenteral nutrition not usually required if isolated
- Antibiotics recommended
- Observe for complications such as wound infection and abscess

Large Intestine







Large Bowel Injury

Incidence

- One of the most lethal abdominal injuries
- Mortality affected by associated injuries
- Penetrating injury is the most common

Management


- Early recognition and control of contamination
- Exploratory laparotomy with primary repair and colostomy
- Preoperative antibiotics
- Observe for complications



Liver Injuries

Liver Functions

- Detoxification
- Synthesis of plasma proteins
- Storage of iron and vitamins
- Metabolism of carbohydrates, protein, and fats
- Phagocytization of bacteria





Liver Injury

Incidence

- Commonly injured organ
- MVC most common cause
- Mortality ~10% to 15%

Liver Injury

Assessment


- Suspect in any patient with blunt injury to right side
- Penetrating trauma produces a range of injuries
- FAST, CT scan
- Grading system




Liver Injury

Management

- Nonoperative management in select patients
- OR for complex lacerations; arterial blush
- Angioembolization
- Aggressive intraoperative resuscitation
- Possible damage control








Liver Injury

Observe for complications

- Recurrent bleeding
- Hemobilia
- Abscess
- Biliary fistula
- Arterial-portal venous fistula
- Sepsis
- Liver failure


Spleen

- Lymphoid organ
- Reservoir for blood
- Vulnerable to injury
- Vascular supply
- Primary immune defense organ





Injuries to the Spleen





Splenic Injury

Incidence

- 2nd most commonly injured abdominal organ
- Mortality depends on the type of trauma and associated injuries
- Mortality related to uncontrolled hemorrhage, delayed rupture, and sepsis



Splenic Injury

Assessment

- Suspect in any patient with blunt injury to left side
- Penetrating trauma can produce a range of injuries
- FAST, CT scan, Angio
- Grading system



Splenic Injury

Management

- Nonoperative in select patients
- Splenorrhaphy and partial splenectomy
- Splenectomy
- Aggressive intraoperative resuscitation
- Possible damage control





Splenic Injury

Management

- Monitor for failed observation
- Observe for postoperative complications
 - Bleeding
 - Thrombocytosis
 - Gastric distention
 - Pancreatitis
 - Infection
- Ensure vaccines are given prior to discharge

Discharge Instructions






Overwhelming Postsplenectomy Sepsis (OPSI)


- Rare
- Can occur from 1 to 5 years after surgery
- Illness presents with flu-like symptoms, shock from sepsis, and DIC followed by death
- Mortality is 50%
- Preventative measures include vaccinations, chemoprophylaxis and education




Retroperitoneal Hemorrhage

- Management depends on the location
- Penetrating trauma requires exploration
- Blunt trauma – pelvic fractures
- Hematoma – explore vs. leave alone






Abdominal Vascular Injuries



Abdominal Vascular Injury

Arterial injury


- Can stop bleeding spontaneously
- Usually occur with pelvic, thoracic, or visceral injury
- Vascular signs may be obscured initially
- Symptoms may include abdominal pain, back pain, hypoactive bowel sounds, tender abdominal mass



Arterial Vascular Injury


Arterial Injury Management

- BP control
- Blood replacement
- Immediate surgery
- End-to-end anastomosis or graft
- Monitor for adequate volume status postoperatively





Venous Vascular Injury




Venous Injury Management

- Quick assessment
- Massive fluid resuscitation
- Pressure and packing
- Operative repair to include ligation and grafting
- Monitor for complications

REBOA

- Resuscitative Endovascular Balloon Occlusion of the Aorta
- Management of non-compressible hemorrhage
- Risk of ischemia and reperfusion injury
- No high grade evidence for improved outcomes



Damage Control



Damage Control

- Abbreviated laparotomy
- Containment of bleeding and contamination
- Temporary intra-abdominal packing
- ICU for physiologic restoration
- Definitive repair





Damage Control

Three phases:

- Control hemorrhage and contamination
- Continued resuscitation in ICU
- Planned reoperation for removal of packing; definitive repair with attempted closure



Did You Know?


- Generally speaking, a retained sponge at any time during the period which the abdomen is “open” with a VacPak™ or Whitman patch™ is not considered an adverse event and is simply part of the management of “Damage Control” and the open abdomen.
- Check own institutional protocols for details and specific policies

Surgical Counts


- X-ray detectable sponges
- Count “incorrect” on operative record
- Obtain an x-ray at end of permanent closure
- Document when x-ray is done in lieu of count



Complications of Abdominal Trauma




Abdominal Compartment Syndrome



Abdominal Compartment Syndrome


Primary Causes:


- Resuscitation edema
- Bowel edema
- Postoperative hemorrhage
- Bowel obstruction
- Closure of abdomen under tension
- Abdominal packing



What Happens to the Body's Organs?

A Vicious Cycle







Reperfusion Phenomenon

- Decompression
 - Release of accumulated acids, metabolites (byproducts of anaerobic metabolism)
 - Profound cardiac depression and hypotension
- To blunt effects
 - 50 mEq Bicarb – up to 4 amps may be ordered
 - Volume resuscitation



Acute Acalculous Cholecystitis (AAC)

- Acute inflammation of gallbladder
- Masked by concomitant injuries and interventions
- Contributing factors include decreased oral intake, TPN, use of narcotics and gallbladder ischemia may occur due to hypotension
- Diagnosis assisted by US, elevated WBC
- Requires surgical intervention



Common Pitfalls

- Failure to suspect intra-abdominal injury from the mechanism of injury
- Failure to fully evaluate abdominal pain after sustaining blunt abdominal injury
- Failure to prepare patient for timely operative intervention
- Failure to recognize hemodynamic compromise and support delay of surgery for additional diagnostic testing



General Nursing Considerations

- Preparation of patient
- Current knowledge of resuscitation
- Administer blood and blood products as ordered
- Prevent hypothermia
- Ongoing monitoring of patients
- Monitor intake and output
- Evidenced-based practice

Summary

- Abdominal trauma presents challenges.
- Not all injuries are easy to diagnose.
- Not all diagnostic modalities are useful in certain injuries.
- Nursing staff must be astute in assessment skills and injury management.
- Teamwork is essential.
- Optimizing outcomes is important.