



SOCIETY OF TRAUMA NURSES



JOINTLY ACCREDITED PROVIDER™  
INTERPROFESSIONAL CONTINUING EDUCATION

# THE ELECTRONIC LIBRARY OF TRAUMA LECTURES

In support of improving patient care, MaineHealth is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team. MaineHealth is also approved to accredit continuing education activities with the following credit types: Interprofessional Continuing Education (IPCE), Association of Social Work Boards (ASWB), American Academy of Physician Assistants (AAPA), American Psychological Association (APA), Registered Dietitians (RDs) and Dietetic Technicians, Registered (DTRs).

THE ELECTRONIC LIBRARY OF

# TRAUMA LECTURES

## Abdominal Trauma





# Objectives

**At the conclusion of this presentation  
the participant will be able to:**

- Describe common mechanisms of injury seen in abdominal trauma
- Discuss various injuries of the abdomen
- State appropriate assessment and diagnostic studies for the patient with abdominal trauma
- Describe abdominal compartment syndrome and the importance of early recognition



# Epidemiology

## Incidence

- Abdominal injuries rank among the top seven causes of death in trauma.
- Accounts for more than 10% of trauma deaths
- Seldom a single system injury

# Mechanism of Injury

**Blunt**



**Penetrating**




By Bobjgalindo - Own work, CC BY-SA 4.0, <https://commons.wikimedia.org/w/index.php?curid=10439138>

# Mechanism of Injury


- Heightens suspicion for certain injuries
- Blunt injury and types of forces
- Use of restraint devices
- Penetrating trauma



# Anatomy and Physiology



# Abdominal Sections



Henry Vandyke Carter, Public domain, via Wikimedia Commons




Image courtesy O.Chaigasame (<http://creativecommons.org/licenses/by/4.0/>)



# Abdominal Assessment


- Inspection
- Auscultation
- Percussion
- Palpation





- Right upper quadrant (RUQ)
- Left upper quadrant (LUQ)
- Right lower quadrant (RLQ)
- Left lower quadrant (LLQ)

# Four Quadrants




## RUQ

- Liver
- Gallbladder with biliary tree
- Duodenum
- Head of pancreas
- Hepatic flexure of colon

## LUQ

- Stomach
- Spleen
- Left lobe liver
- Left kidney
- Left adrenal gland
- Splenic flexure of colon
- Parts of transverse and descending colon





## RLQ

- Cecum
- Appendix
- Ascending colon
- Right ovary and fallopian tube
- Right ureter

## LLQ

- Descending colon
- Sigmoid colon
- Left ovary and fallopian tube
- Left uterine tube





# Ongoing Assessment

- Delayed diagnosis or missed injuries
- Frequent serial and systematic examinations
- Tertiary exam

# Diagnostic Labs

Are they necessary?

Reliable?

- Hematocrit
- WBC
- Electrolytes



# Diagnostic Labs

- ABGs
- Coagulation studies
- Urinalysis
- Pregnancy
- Pancreatic enzymes
- LFTs



# Diagnostic Modalities

- Radiographs
- Diagnostic peritoneal lavage (DPL)
- Ultrasonography (US)
- Computed tomography (CT) scan
- Angiogram
- Diagnostic laparoscopy






# Radiographic Films

- CXR
  - Concomitant pulmonary and cardiac injuries
  - Displacement of abdominal organs
- Pelvis
- Plain abdominal films have limited if any role in the acute resuscitation.
  - AP and lateral films may identify fluid or air.
  - Upright film for free air; may disclose ruptured hollow viscus

# Diagnostic Peritoneal Lavage

- Used to diagnose intra-abdominal bleeding
- Indications
  - Unexplained hypotension, decreased hematocrit, or shock
  - CT or ultrasound not available
  - Equivocal abdominal examination
  - Altered mental status
  - Spinal cord injury
  - Distracting injuries



# DPL

## Advantages

- Quick, simple
- Safe
- Low cost
- Relatively accurate
- Grossly positive result

## Disadvantages

- Difficult to perform in some patients
- Invasive procedure
- Can miss certain injuries

**Complications:** Infection, hematoma, false positives, injury, bleeding, unnecessary laparotomy, failure to recover lavage fluid

**Note: A urinary catheter and gastric tube should be in place prior to the procedure.**



# Ultrasound

## FAST

- Focused
- Assessment
- Sonography
- Trauma

## Ultrasound probe locations and sequence

- Epigastrium
- RUQ
- LUQ
- Pelvis



# Ultrasound

- Reliable, fast, safe
- Noninvasive
- Equipment portable
- Performed simultaneously
- Fast exam detects free fluid
- Serial exams
- Leads to fewer DPL's & CT Scans





# Disadvantages of Ultrasound

- Clinician expertise variable
- Lacks specificity & sensitivity
- Reliability is questionable
- May not reveal free fluid if performed too early


# Computed Tomography



- Used for hemodynamically stable patients
- Advantages:
  - Noninvasive procedure
  - Better defines organ injury
  - Estimates amount of blood in spaces
  - Retroperitoneum and vertebrae can be assessed
- Helical scanners

# CT Scan in Trauma

- Visualizes abdominal solid organs and vessels well
- Does NOT see mesenteric injuries, hollow viscus, duodenum, diaphragm, or omentum well
- Whole body scans on all trauma
- Radiation long-term effects



# CT Scan Disadvantages

- Takes time to perform
- Cost
- Transport of patient
- Requires stable and cooperative patient
- Less reliable in diagnosing some injuries
- IV contrast
- Radiation exposure





# Angiography

- Detects active bleeding in patients with vascular trauma
- Embolizes specific structures within bleeding organs or the pelvis
- Detects A-V fistulas and aneurysms in penetrating trauma



# Diagnostic Laparoscopy (DL)

- Screening or diagnostic tool
- Invasive procedure with some limitations
- Used to detect or exclude certain findings
- May reduce the rate of negative laparotomies





# Other Diagnostic Procedures

## ERCP

- May be indicated in the stable trauma patient suspected of having biliary tract or pancreatic duct injury
- Most accurate test in the patient with hyperamylasemia and in those following pancreatic surgery

# Other Diagnostic Procedures

## Gastrografin or barium studies

- Helpful in diagnosing injuries to the esophagus, stomach, or bowel
- Contrast enemas are used to diagnose rectal or colonic injury secondary to penetrating trauma





Image courtesy S Bhimji MD, <https://www.statpearls.com/ArticleLibrary/viewarticle/21348>

# Specific Injuries





# Esophageal Injuries



# Esophagus

## Anatomy

- Carries food from pharynx to the stomach
- Joins the stomach at the level of T-10
- Posterior surface overlies aorta
- Anterior surface covered by peritoneum

## Predisposing Injury Facts

- Narrow at
  - Cricoid cartilage
  - Arch of aorta
  - Esophagogastric junction
- Lacks serosal layer
  - Integrity of anastomoses
  - Possible leak after surgical repair



# Esophageal Injury

- Incidence
  - Higher in cervical and thoracic areas
  - Majority are due to penetrating trauma
  - Blunt injury is rare
- Early diagnosis essential
- Can result in high morbidity and mortality

## Sequelae

- Respiratory compromise
- Mediastinitis
- Paraesophageal abscess
- Empyema
- Esophageal fistula
- Peritonitis



# Esophageal Injury

## Assessment

- Symptoms of perforation include pain, fever, and dysphagia
- Symptoms of abdominal esophageal tear include signs of peritoneal irritation followed by dyspnea and pleuritic pain

## Diagnostic tests

- Endoscopy/Esophagoscopy
- CT



# Esophageal Injury




## Management

- Initial assessment complex
- Goal is to minimize the bacterial contamination and enzyme erosion
- Gastric decompression
- Antibiotic coverage
- Drainage of wound
- Surgical repair

# Esophageal Injury Management

Continuous monitoring for injury

Complications after repair



Samir धर्म, Public domain, via Wikimedia Commons



# Diaphragm



# Diaphragmatic Injury

## Incidence

- Usually occurs with other injuries
- Seen in < 5% of blunt trauma patients
- Left side greater incidence than right side
- Commonly associated with penetrating trauma
- Injuries from blunt trauma caused by sudden rise in intrathoracic pressure



# Diaphragmatic Injury


## Assessment


- Auscultation of peristaltic sounds in chest
- Delayed rupture - unexplained chest pain and increased respiratory rate
- CXR is most important diagnostic study
  - Elevation of hemidiaphragm
  - Bowel pattern in the chest
  - Gastric tube curls in chest
  - Hemothorax – associated injury
- Masked by positive pressure ventilation

TRAUMA

SUPINE

L  
T83








# Diaphragmatic Injury


## Management

- Exploratory laparotomy
- Diagnostic laparoscopy in penetrating trauma










# Pancreatic Injuries







# Pancreatic Injuries

## Incidence

- Uncommon (0.2% to 12%)
- Associated with other abdominal injuries
- Majority caused by penetrating trauma
- Blunt trauma is usually a direct blow or compression type force
- Mortality is variable
- Pancreatic Injuries graded by severity I-V
- Most grades of III or higher will require OR




# Duodenum



# Duodenal Injuries

## Incidence


- Majority caused by penetrating trauma
- Blunt trauma is usually compression type
- Mortality is variable
- Multi-organ injuries



# Pancreatic and Duodenal Injuries

## Assessment

- Peritoneal symptoms not evident but appear later
- CT scan is the exam of choice
- Injury usually found intraoperatively





# Pancreatic and Duodenal Injuries

## Assessment


- Blunt injury to duodenum can produce intramural hematoma
- Perforation causes contamination



# Pancreatic and Duodenal Injuries

## Management

- Options depend on site and severity
- Primary closure
- Simple external closed drainage
- Distal pancreatectomy
- Pancreatic duodenectomy





# Pancreatic and Duodenal Injuries

## Duodenal Management

- Debridement and primary repair
- Surgical procedure depends on hemodynamic stability and duct involvement
- Nonoperative management requires close observation

## Pancreatic Management

- Primary cause of death is hemorrhage
- Late deaths are due to sepsis, ARDS, multiple organ failure
- Observe for complications

An intraoperative photograph showing a surgical incision in the abdominal wall. The incision is a vertical laceration with jagged edges, exposing the underlying muscle and peritoneum. A significant amount of bright red blood is visible, pooling in the wound and on the surrounding tissues. Several gloved hands are visible, holding back muscle and soft tissue to provide a clear view of the surgical site. The lighting is bright, highlighting the moist, glistening surfaces of the internal organs and the texture of the muscle and skin.

# Injuries to the Stomach and Intestines

# Stomach Injury

## Incidence

- Rare; more common in children
- Commonly involves adjacent organs
- Protected by location and mobility
- Most common cause is penetrating injury





# Stomach Injury

## Assessment

- Symptoms variable and nonspecific
- May include severe epigastric or abdominal pain, tenderness, signs of peritonitis
- Clouded by associated injuries
- Bloody output from NG tube
- Free air on radiograph
- Findings on CT or DPL



# Stomach Injury

- Indications for surgery
  - Hemodynamic instability
  - Diffuse abdominal pain/peritonitis on clinical exam
  - Studies consistent with GI perforation
- Selective non-operative management
  - Stable
  - No peritoneal signs
  - No clear radiographic evidence




# Stomach Injury

## Management

- Gastric decompression
- Decision for surgical intervention vs. non-operative management
- If contamination exists, copious peritoneal irrigation and delayed primary closure
- Monitor for postoperative complications



# Intestinal Injuries





# Small Intestine

## Jejunum and Ileum

- Responsible for nutrient absorption and fluid and electrolyte shifts
- Jejunum lies in the umbilical region
- Ileum lies in the hypogastric and pelvic regions
- Vulnerable to seatbelt injury





# Small Bowel Injury

## Incidence

- Most frequently injured by penetrating trauma.
- Blunt injury is relatively uncommon.
- Presence of pancreatic and solid organ injury are predictive of increased risk for hollow viscus injury.

## Assessment

- Clinical signs may not be apparent initially.
- Blunt vs penetrating
- Signs of peritonitis develop.
- Any blow to the abdomen or penetrating injury to the lower chest or abdomen should increase suspicion of injury.





# Small Bowel Injury

## Management

- Bleeding should be controlled prior to exploration
- Debridement followed by primary closure and ligation of bleeders
- Bowel resection for multiple defects
- Gastric decompression and parenteral nutrition not usually required if isolated
- Antibiotics recommended
- Observe for complications such as wound infection and abscess

# Large Intestine







# Large Bowel Injury

## Incidence

- One of the most lethal abdominal injuries
- Mortality affected by associated injuries
- Penetrating injury is the most common

## Management


- Early recognition and control of contamination
- Exploratory laparotomy with primary repair and colostomy
- Preoperative antibiotics
- Observe for complications




# Liver Injuries

# Liver Functions

- Detoxification
- Synthesis of plasma proteins
- Storage of iron and vitamins
- Metabolism of carbohydrates, protein, and fats
- Phagocytization of bacteria





# Liver Injury

## Incidence

- Commonly injured organ
- MVC most common cause
- Mortality ~10% to 15%

# Liver Injury

## Assessment


- Suspect in any patient with blunt injury to right side
- Penetrating trauma produces a range of injuries
- FAST, CT scan
- Grading system




# Liver Injury

## Management

- Nonoperative management in select patients
- OR for complex lacerations; arterial blush
- Angioembolization
- Aggressive intraoperative resuscitation
- Possible damage control








# Liver Injury


## Observe for complications

- Recurrent bleeding
- Hemobilia
- Abscess
- Biliary fistula
- Arterial-portal venous fistula
- Sepsis
- Liver failure

# Spleen

- Lymphoid organ
- Reservoir for blood
- Vulnerable to injury
- Vascular supply
- Primary immune defense organ





# Injuries to the Spleen



# Splenic Injury

## Incidence

- 2<sup>nd</sup> most commonly injured abdominal organ
- Mortality depends on the type of trauma and associated injuries
- Mortality related to uncontrolled hemorrhage, delayed rupture, and sepsis



# Splenic Injury

## Assessment

- Suspect in any patient with blunt injury to left side
- Penetrating trauma can produce a range of injuries
- FAST, CT scan, Angio
- Grading system

# Splenic Injury

## Management

- Nonoperative in select patients
- Splenorrhaphy and partial splenectomy
- Splenectomy
- Aggressive intraoperative resuscitation
- Possible damage control





# Splenic Injury

## Management

- Monitor for failed observation
- Observe for postoperative complications
  - Bleeding
  - Thrombocytosis
  - Gastric distention
  - Pancreatitis
  - Infection
- Ensure vaccines are given prior to discharge

# Discharge Instructions






# Overwhelming Postsplenectomy Sepsis (OPSI)


- Rare
- Can occur from 1 to 5 years after surgery
- Illness presents with flu-like symptoms, shock from sepsis, and DIC followed by death
- Mortality is 50%
- Preventative measures include vaccinations, chemoprophylaxis and education




# Retroperitoneal Hemorrhage

- Management depends on the location
- Penetrating trauma requires exploration
- Blunt trauma – pelvic fractures
- Hematoma – explore vs. leave alone






# Abdominal Vascular Injuries



# Abdominal Vascular Injury

## Arterial injury


- Can stop bleeding spontaneously
- Usually occur with pelvic, thoracic, or visceral injury
- Vascular signs may be obscured initially
- Symptoms may include abdominal pain, back pain, hypoactive bowel sounds, tender abdominal mass



# Arterial Vascular Injury


## Arterial Injury Management

- BP control
- Blood replacement
- Immediate surgery
- End-to-end anastomosis or graft
- Monitor for adequate volume status postoperatively





# Venous Vascular Injury




## Venous Injury Management

- Quick assessment
- Massive fluid resuscitation
- Pressure and packing
- Operative repair to include ligation and grafting
- Monitor for complications

# REBOA

- Resuscitative Endovascular Balloon Occlusion of the Aorta
- Management of non-compressible hemorrhage
- Risk of ischemia and reperfusion injury
- No high grade evidence for improved outcomes



# Damage Control



# Damage Control

- Abbreviated laparotomy
- Containment of bleeding and contamination
- Temporary intra-abdominal packing
- ICU for physiologic restoration
- Definitive repair





# Damage Control

## Three phases:

- Control hemorrhage and contamination
- Continued resuscitation in ICU
- Planned reoperation for removal of packing; definitive repair with attempted closure



# Did You Know?

- Generally speaking, a retained sponge at any time during the period which the abdomen is “open” with a VacPak™ or Whitman patch™ is not considered an adverse event and is simply part of the management of “Damage Control” and the open abdomen.
- Check own institutional protocols for details and specific policies

# Surgical Counts


- X-ray detectable sponges
- Count “incorrect” on operative record
- Obtain an x-ray at end of permanent closure
- Document when x-ray is done in lieu of count



# Complications of Abdominal Trauma




# Abdominal Compartment Syndrome



# Abdominal Compartment Syndrome


## Primary Causes:


- Resuscitation edema
- Bowel edema
- Postoperative hemorrhage
- Bowel obstruction
- Closure of abdomen under tension
- Abdominal packing



# What Happens to the Body's Organs?

## A Vicious Cycle





# Reperfusion Phenomenon

- Decompression
  - Release of accumulated acids, metabolites (byproducts of anaerobic metabolism)
  - Profound cardiac depression and hypotension
- To blunt effects
  - 50 mEq Bicarb – up to 4 amps may be ordered
  - Volume resuscitation

# Acute Acalculous Cholecystitis (AAC)

- Acute inflammation of gallbladder
- Masked by concomitant injuries and interventions
- Contributing factors include decreased oral intake, TPN, use of narcotics and gallbladder ischemia may occur due to hypotension
- Diagnosis assisted by US, elevated WBC
- Requires surgical intervention



# Common Pitfalls

- Failure to suspect intra-abdominal injury from the mechanism of injury
- Failure to fully evaluate abdominal pain after sustaining blunt abdominal injury
- Failure to prepare patient for timely operative intervention
- Failure to recognize hemodynamic compromise and support delay of surgery for additional diagnostic testing



# General Nursing Considerations

- Preparation of patient
- Current knowledge of resuscitation
- Administer blood and blood products as ordered
- Prevent hypothermia
- Ongoing monitoring of patients
- Monitor intake and output
- Evidenced-based practice

# Summary

- Abdominal trauma presents challenges.
- Not all injuries are easy to diagnose.
- Not all diagnostic modalities are useful in certain injuries.
- Nursing staff must be astute in assessment skills and injury management.
- Teamwork is essential.
- Optimizing outcomes is important.