



SOCIETY OF TRAUMA NURSES



JOINTLY ACCREDITED PROVIDER™

INTERPROFESSIONAL CONTINUING EDUCATION

THE ELECTRONIC LIBRARY OF

# TRAUMA LECTURES

In support of improving patient care, MaineHealth is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

MaineHealth is also approved to accredit continuing education activities with the following credit types: Interprofessional Continuing Education (IPCE), Association of Social Work Boards (ASWB), American Academy of Physician Assistants (AAPA), American Psychological Association (APA), Registered Dietitians (RDs) and Dietetic Technicians, Registered (DTRs).

THE ELECTRONIC LIBRARY OF

# TRAUMA LECTURES

## Obesity and Trauma



SOCIETY OF TRAUMA NURSES

# Objectives

**At the conclusion of this presentation,  
the participant will be able to:**

- Describe how the obesity epidemic impacts the delivery of trauma care
- Describe how obesity impacts body systems
- Discuss the challenges and considerations associated with resuscitation and management of the obese trauma patient
- Describe management approaches to care of the injured obese patient with blunt, penetrating, or burn injuries


# Introduction



1 out of every 3 U.S.  
adults is obese.  
(Harvard, 2020)



# Body Mass Index





# Consequences of Obesity

- Obesity is associated with the leading causes of death in the United States and worldwide, including:
  - Diabetes
  - Heart disease
  - Stroke
  - A wide range of cancers
- Low quality of life
- Mental illness such as clinical depression, anxiety, and other mental disorders
- Body pain and difficulty with physical functioning



# Societal Costs

## Direct

- 2016, the aggregate in medical cost due to obesity among adults in the United States was \$260.6 billion. (Cowley, 2021)
- The effects of obesity raised costs in every category of care: inpatient, outpatient, and prescription drugs.

## Indirect


- Absent from work for obesity-related health reasons
- Decreased productivity while at work
- Premature death and disability



# Injury Patterns


- Increased BMI = increased rates
  - Extremity fractures (humerus, femur, tibia/fibula)
  - Chest injuries
  - Spinal injuries
  - Increased mortality
  - Increased complications
- Increased BMI = decreased rates
  - Hip fractures
  - Head injuries (more fatal)
  - Liver lacerations

## Trends in obesity among children and adolescents aged 2–19 years, by age: United States, 1963–1965 through 2013–2014



NOTES: Obesity is defined as body mass index (BMI) greater than or equal to the 95th percentile from the sex-specific BMI-for-age 2000 CDC Growth Charts.

SOURCES: NCHS, National Health Examination Surveys II (ages 6–11) and III (ages 12–17); and National Health and Nutrition Examination Surveys (NHANES) I–III, and NHANES 1999–2000, 2001–2002, 2003–2004, 2005–2006, 2007–2008, 2009–2010, 2011–2012, and 2013–2014.



# Cardiovascular


- Obesity is an independent risk factor for cardiac disease.
- Increased circulating blood volume, cardiac output
- Increased risk of thromboembolism
- Increased systemic vascular resistance





# Pulmonary

- Increased chest wall resistance
- Increased intra-abdominal pressure
- Dysfunctional chest wall
- Obstructive sleep apnea




Total protein	
Albumin	
Globulin	
Total bilirubin	
Direct bilirubin	
AST (SGOT)	220 H
ALT (SGPT)	340 H
ALP	
GGT	

## GI/Liver/Renal

- Decreased gastric motility
- Increased intra-abdominal pressure
- Gastroesophageal reflux
- Increased aspiration risk during intubation
- Chronic elevations liver enzymes
- Pancreatitis
- Chronic renal failure

# Endocrine/Immunologic


- Inflammation
- Metabolic syndrome
- Elevated cortisol levels



# Musculoskeletal

- Normal increased stress on bones and joints
  - Can be described as severe pain
- Osteoarthritis
- Osteoporosis



A person wearing a light-colored short-sleeved shirt and dark pants is riding a black bicycle on a city street. The person is holding a white water bottle in their left hand. The background shows a blurred city street with other bicycles and a blue vehicle. A semi-transparent white box is overlaid on the left side of the image, containing text.

**From the field, prehospital, to the acute care setting, obstacles arise that complicate and delay the care of the obese and morbidly obese patient.**



# Pre-Hospital

- Prolonged extrication time
- O2 masks may not fit
- Larger C Collars
- Inability to lie flat
- Stretchers/backboards limited size and loads
- Increased staff
- Assistive equipment
- Evacuation
- BP cuffs



The BEAR stair chair, Beariatrics.com

# Emergency Department

- Monitoring
  - BP cuffs
  - Pulse ox
  - Introducers



- Supplies
  - Stretcher
  - Wheelchairs
  - Splints
  - Tourniquets
  - Scales
- Diagnostics
  - Radiology tables, plates
  - CT/MRI tables, circumference
  - OR
- Staffing ratio



# Admitted Patient Care Rooms

- Bed
- Portable or fixed overhead lifts
- Scales
- Extra large BP cuffs
- Wheelchairs
- Toilets/shower/grab bars
- Bedside commode
- Gowns/bottoms

# Assessment Principles

- Primary
- Secondary
- Disposition
- Management





# Airway (C-Spine Protection)

## CHALLENGES

- Short thick necks
- Poor extension
- Loss of landmarks
- Adipose tissue
- Fat deposits in pharyngeal tissue
- Gastro-esophageal reflux
- Backboard weight limits



# Airway (C-Spine Protection)

## CONSIDERATIONS

- Position with head of bed slightly elevated or reverse Trendelenburg
- Use of sandbags and tape for immobilization
- Gastric tube insertion if concern for reflux/emesis/airway
- Dedicated member to maintain c-spine control
- Early surgical cricothyrotomy
- Optical equipment (i.e. video laryngoscope)
- History of gastric banding

# Breathing

## CHALLENGES

- Fat deposits in diaphragm and intercostal muscles
- Elevated diaphragm
- Rapid desaturation
- Chest weight
- Increased work of breathing
- Sleep apnea
- Impaired lung compliance
- Difficulty auscultating breath sounds







# Breathing

## CONSIDERATIONS

- CPAP/PEEP
- Reverse Trendelenburg
- 2-person bag-mask
- Needle decompression/chest tube placement
- “Awake” intubation vs RSI/DAI
- Longer recovery time with failed attempted intubation
- Neck circumference
- Ramping position





# Circulation

## CHALLENGES

- Difficulty palpating central and peripheral pulses
- BP assessment
- Auscultation heart sounds
- High oxygen supply and demand
- IV access
- Accuracy of pulse ox






# Circulation

## CONSIDERATIONS

- Longer introducers
- Use of intra-osseous needle
- Arterial line
- Intra-abdominal hypertension
- Lactate
- pH
- CVP
- PA catheter
- TEE





# Disability

## Considerations

- CPAP at home?
- Lack of mobility
- Toileting

## Challenges

- Establish baseline
- Early discharge planning



# Exposure/Environment

- Skin shearing
- Hypothermia
- Inspect for skin rashes, fungal infections, decubitus, wounds
- Large pannus
- Larger patient gowns
- Moving boards
- Assistance
- Stretchers/ beds



# Secondary Survey

## CONSIDERATIONS

- Clinical exams less reliable
- Skin folds may mask penetrating injury
- Difficult to assess abdominal or bone tenderness
- Masses/deformities difficult to palpate
- Use of inappropriate cuff size
- Difficult to assess the back
- Imaging issues





# Give Comfort

## CHALLENGES

- Patient Size
- Bias
- Stigma
- Psychosocial Issues

## CONSIDERATIONS

- Addressing bias may be first step to improving outcomes
- Medications
  - Right dose
  - Right route
- Specialized beds and equipment



# History/Inspect Posterior

## History

- Pre-hospital
- Referring Facility
- Medications
- Co-morbidities
- Surgeries

## Inspect the Back

- Number of people needed to log roll
- Patient safety
- Bed width
- Skin folds



<https://www.healthcaredailyonline.com/hospitals-using-obesity-suit-to-better-understand-patients/>

# Disposition

- Decide early if transfer is necessary.
- Inform transferring agency of patient's size.
- Inform admitting unit of the patient's size ASAP to allow them to prepare.
- Make preparations to ensure patient safety.
  - Interfacility
  - Intrafacility



# Nutritional Requirements

- Nutritional requirements differ
- Increased caloric requirements in trauma, but hypocaloric high protein diet preferred in obese patients
- High percentage of Vitamin D deficiency in obesity
- May need indirect calorimetry to prescribe diet
- Maintain blood sugar control
- Monitor weight



Global Industrial™ Wheelchair Scale, 1,000 lb x 0.5 lb



# Pharmacology

- Consult on admission for your PharmD or Clinical Pharmacist to review all current medications and to compute dosing of:
  - Antibiotics
  - Anti-thrombotics
  - Analgesics
- Restart home medications when appropriate
- Individualize pain management

- Calculations

- Dose weight (DW)
- Ideal body weight (IBW)
- Total body weight (TBW)

$$DW = IBW + 0.3 (TBW - IBW)$$

# Labs to Watch

- BUN
- Creatinine
- Insulin levels
- Cortisol
- Hyperlipidemia
- PT/PTT
- C-reactive protein
- Cytokine
- Lactate
- Amylase/lipase
- ABG
- Anti-Xa
- Growth hormone
- Prolactin
- TSH



# Missed Injury/Delayed Diagnosis

- Sternal fractures
- Flail chest
- Pelvic fractures
- Rib fractures
- Pulmonary contusions
- Long bone fractures

Limitations with traditional evaluations:

- Chest x-ray
- Ortho x-rays
- FAST
- CT scan
- Exploratory laparotomy



# Complications

- Atelectasis
- Anastomotic leaks
- Higher risk of re-operation
- Higher infection rates
- Abdominal compartment syndrome
- Thrombophlebitis/PE
- Urinary tract infections
- Decubiti



# Fractures

- Strength of rods
- Compartment syndrome
- Casting more difficult
- Vascular assessment challenges
- Higher rate of amputation
- Vitamin D insufficiency
- During hip and knee arthroplasty, the infection rate is nearly 5% in obese patients and nearly 10% in obese, diabetic patients.





# Consultations

- Nutrition
  - Hypocaloric
  - Higher protein
  - Prebiotics/probiotics
- Pharm D
- Primary care providers
- Case management
- Social work
- Sleep apnea



# Functional Independence Measurement (FIM)

- Admission, discharge, 6 months post-discharge
- Domains
  - Self-care
  - Sphincter control
  - Mobility
  - Locomotion
  - Communication
  - Social cognition
- Compared with nonobese patients, the rate of recovery was reduced by 30% in overweight, 37% in obese, and 48% in morbidly obese patients. (Dhungel 2015)



# Mortality Risk

- There was no statistically significant difference in mortality. (Drury 2021)
- Severely obese trauma patients were at least 30% more likely to die and approximately twice as likely to have a major complication. (Glance 2014)
- Obese trauma patients undergoing emergent trauma laparotomy have a high likelihood for both complications and mortality, with morbidly obese trauma patients having the highest likelihood for both. (Covarrubias 2021)
- In a cohort of matched patients, morbid obesity is a risk factor for the development of in-hospital complications and mortality after blunt traumatic injury. (Ditillo 2014)
- Increasing BMI by category was associated with a stepwise increase in odds of acute kidney injury, cardiovascular events, total hospital length of stay (LOS), intensive care unit LOS, and ventilator days. (Hakam 2021)

# Summary

- Obesity places challenges on healthcare costs by stressing infrastructure, requiring specialized equipment, and evidence-based education.
- Obesity independently impacts body systems, thus increasing the complexity of trauma care.
- Anticipatory preparation for the challenges and barriers to providing good trauma care to the obese patient will lead to provision of optimum care.